

# Influenza A virus transmission: contributing factors and clinical implications

Jessica A. Belser, Taronna R. Maines, Terrence M. Tumpey  
and Jacqueline M. Katz\*

Efficient human-to-human transmission is a necessary property for the generation of a pandemic influenza virus. To date, only influenza A viruses within the H1–H3 subtypes have achieved this capacity. However, sporadic cases of severe disease in individuals following infection with avian influenza A viruses over the past decade, and the emergence of a pandemic H1N1 swine-origin virus in 2009, underscore the need to better understand how influenza viruses acquire the ability to transmit efficiently. In this review, we discuss the biological constraints and molecular features known to affect virus transmissibility to and among humans. Factors influencing the behaviour of aerosols in the environment are described, and the mammalian models used to study virus transmission are presented. Recent progress in understanding the molecular determinants that confer efficient transmission has identified crucial roles for the haemagglutinin and polymerase proteins; nevertheless, influenza virus transmission remains a polygenic trait that is not completely understood. The clinical implications of this research, including methods currently under investigation to mitigate influenza virus human-to-human transmission, are discussed. A better understanding of the viral determinants necessary for efficient transmission will allow us to identify avian influenza viruses with pandemic potential.

Influenza A viruses are single-stranded, negative-sense, enveloped RNA viruses, possessing eight gene segments that encode 11 known proteins (Refs 1, 2). The two major surface glycoproteins haemagglutinin (HA) and neuraminidase (NA) form the basis of serologically distinct

virus subtypes. There are currently 16 HA and nine NA subtypes identified; all subtypes are found in wild waterbirds, the natural reservoir for influenza A viruses (Refs 2, 3). Despite this extensive diversity, only select subtypes have breached the species barrier, and, to date,

Influenza Division, National Center for Immunization and Respiratory Diseases, Centers for Disease Control and Prevention, Atlanta, GA, USA.

\*Corresponding author: Jacqueline M. Katz, Influenza Division MS G-16, 1600 Clifton Road NE, Atlanta, GA 30333, USA. E-mail: jkatz@cdc.gov

viruses bearing H1, H2, H3, H5, H7 or H9 have caused documented human infections (Refs 2, 4, 5). The generation of a pandemic requires the emergence of a virus in the human population that (1) bears an HA to which there is little or no pre-existing immunity, (2) causes disease in humans, and (3) is capable of sustained human-to-human transmission (Fig. 1). However, only three subtypes (H1–H3) have acquired all three of these properties to cause pandemics in the last 100 years – in 1918 (H1N1), 1957 (H2N2), 1968 (H3N2) and 2009 (H1N1). In the relatively rare transmission of avian viruses to humans, primarily through contact with infected domestic poultry, H5, H7 and H9 subtypes have failed to establish a foothold in human populations. Therefore, understanding the ability of a virus to achieve a transmissible phenotype is an essential component in assessing the pandemic potential of novel subtypes. In this review, we highlight the complexities inherent in the study of influenza virus transmissibility, how the use of mammalian models and reverse genetics has facilitated a more detailed examination of the molecular determinants that govern virus transmission, and the clinical implications of this knowledge.

### Assessment of influenza A virus transmissibility


The ability of influenza viruses to spread through susceptible populations to cause annual epidemics and occasional pandemics is well documented. When modelling influenza virus transmission in a laboratory setting, accurately defining the route of transmission under investigation is crucial. Influenza viruses spread either by inhalation of virus-containing respiratory droplets or droplet nuclei expelled from an infected person during breathing, coughing, sneezing or talking, or by direct or indirect contact with virus-containing fomites on environmental surfaces (Refs 6, 7, 8). The use of mammalian models has enabled detailed study of influenza virus transmission by both contact and respiratory droplet routes.

### Transmissibility of human and avian influenza A viruses

Historical records suggest frequent appearances of a highly contagious, acute respiratory illness infecting humans since ancient times, with an aetiology consistent with present-day influenza

epidemics (Ref. 9). The efficient transmissibility of seasonal and pandemic influenza viruses is well known; the contagious nature of influenza has been documented by epidemiological studies from outbreaks occurring among diverse populations and locations, including cruise ships, school dormitories, hospitals and nursing homes (Refs 10, 11, 12, 13, 14). The presence of a single ill individual in a confined space, such as an airplane, was sufficient for an outbreak to occur among previously healthy passengers (Refs 15, 16). Analysis of global airline flight itineraries at the onset of the 2009 H1N1 pandemic found a strong correlation between international movement of travellers and countries with confirmed influenza infections, further illustrating the capacity for rapid spread of infectious disease in a highly mobile global population (Ref. 17). The rates of influenza virus infections are highest during pandemics, when little to no pre-existing immunity to the virus exists among susceptible populations, although attack and case-fatality rates vary from pandemic to pandemic (Ref. 18). Nevertheless, once pandemic viruses become established in humans, their efficient seasonal spread among otherwise healthy individuals ultimately provides an ongoing and even greater public health burden in terms of hospitalisations and lives lost. Serious illness and death following seasonal influenza virus infection in the USA occurs predominantly among the elderly, resulting in an average of 36 000 deaths each year (Ref. 19). As such, influenza viruses present not only an occasional pandemic threat but also an ongoing annual threat to public health worldwide.

Unlike human influenza viruses, wholly avian low-pathogenic influenza viruses do not generally replicate to high titres in humans and have not been demonstrated to readily transmit between people (Refs 20, 21). However, sporadic human infections with avian influenza viruses of the H5, H7 and H9 subtypes have occurred, frequently following exposure to infected poultry (Refs 4, 5, 22). Probable human-to-human transmission of highly pathogenic avian influenza (HPAI) viruses of the H5N1 and H7N7 subtypes has been reported rarely (Refs 23, 24, 25, 26). Although these avian influenza viruses have not demonstrated the capacity for sustained transmission in a human population, continual evolution of these viruses in avian species calls for close surveillance of



**Figure 1. Assessing the pandemic potential of influenza viruses.** Of the three properties an influenza virus must possess to cause a pandemic (blue boxes), avian influenza viruses currently meet two (red boxes). Several barriers have been proposed to contribute to the poor transmissibility of avian influenza viruses in humans (green box), but the potential for these viruses to overcome these barriers maintains their status as a pandemic threat (amber box). Abbreviations: HA, haemagglutinin; PB2, polymerase basic protein 2; SA, sialic acid; URT, upper respiratory tract.

land-based poultry and individuals exposed to potentially infected birds for the emergence of a novel strain that might possess or readily acquire this ability. Transmission of influenza viruses between avian species, and transmission and subsequent establishment of influenza virus lineages in other mammals (including swine, equine and canine species), are a threat to public health but fall outside the scope of this review.

### Modes of influenza virus transmission

The diversity of virus-containing particles expelled by infected persons can result in the transmission of influenza viruses to susceptible

hosts by distinct routes (Ref. 27). The infectivity of airborne virus in small respiratory droplets (approximately  $<5 \mu\text{m}$  in diameter) can be very high: the infectious dose of influenza virus in humans following aerosol inhalation was reported to be as low as 3 50% tissue culture infectious doses (TCID<sub>50</sub>) in one study performed with experimentally infected persons (Ref. 6). Transmission of large respiratory droplets (approximately  $>5 \mu\text{m}$  in diameter) can also occur by inhalation of virus particles by a susceptible person, or can contribute to contamination of proximal environmental surfaces. In the absence of respiratory droplets,

transmission can occur by direct physical contact between an infected and susceptible person, or by indirect contact with contaminated environmental surfaces or fomites. Influenza A viruses can survive on hard, nonporous surfaces for 12–24 h, and on tissues, cloth and paper for <8–12 h; transfer of virus from these surfaces to hands can therefore take place up to several hours after initial virus deposition (Ref. 7). Transmission of a seasonal H3N2 influenza virus was found to be more efficient by the aerosol route compared with contact with contaminated environmental surfaces (fomites) in the guinea pig model (Ref. 28). Despite these studies, the relative contribution of contact, droplet or airborne transmission of influenza viruses in humans remains unknown.

Although avian viruses of the H5, H7 and H9 subtype do not currently possess the ability of sustained human-to-human transmission, the hundreds of documented cases of human disease caused by viruses within these subtypes demonstrate the ability of influenza A viruses to infect humans with varying efficiencies and by different routes. The majority of human cases of avian H5 and H7 virus infections have been associated with direct contact transmission from exposure to sick or dead poultry, often following slaughtering and preparing sick poultry for cooking (Ref. 4). For a minority of cases of H5N1 transmission, the only identified risk factor was visiting a live-poultry market, suggesting that one transmission route might be inhalation of aerosolised infectious poultry faeces or other material (Refs 4, 29). Human H5N1 virus infection has also been documented following the consumption of uncooked blood from H5N1-infected ducks (Ref. 4). Experimentally, the consumption of raw H5N1-virus-infected domestic poultry by ferrets, or of virus-infected chicks by cats, was shown to be sufficient to cause infection of the respiratory tract in addition to the digestive tract (Refs 30, 31). Such data suggest that the respiratory tract can be infected by the oral consumption of uncooked poultry products, possibly through direct pharyngeal or tonsillar infection or inhalation of aerosolised virus during mastication. For select avian influenza viruses of the H7 subtype, transmission to humans involved initial conjunctival infections that in isolated cases have presented with respiratory disease (Refs 23, 32, 33). Thus, the conjunctiva might represent a primary

portal of entry for H7 viruses. These ongoing human infections with viruses that do not currently have the ability to readily transmit among humans nonetheless constitute a public health threat and highlight the importance of understanding the mechanisms used by influenza A viruses to transmit by contact or airborne routes.

### **Mammalian models of influenza virus transmission**

The ferret has been used to model influenza infection since it was first used to isolate the virus in the 1930s. Ferrets are naturally susceptible to influenza viruses and exhibit many clinical signs following infection (such as sneezing, fever and nasal discharge) that are comparable to influenza-like illness in humans (Refs 34, 35). Furthermore, the respiratory tract of ferrets appears to closely resemble that of humans, because avian H5N1 and human H3N2 influenza viruses display similar patterns of virus attachment to respiratory tissues of both species (Refs 36, 37). The utility of ferrets to assess the transmissibility of influenza viruses was described as early as 1941 in a study by Andrewes and Glover, who found that susceptible naive ferrets housed in the same room as ferrets shedding virus became infected, even when physical barriers that would trap and prevent the dissemination of large droplets were used (Ref. 38). The ferret model has in recent years become the model of choice for most influenza virus research and is routinely used to examine the transmissibility of both human and avian influenza viruses by either direct contact or respiratory droplets (Refs 39, 40, 41, 42, 43). The high susceptibility of ferrets to influenza virus infection can make it difficult to obtain ferrets that are seronegative against currently circulating influenza viruses for research purposes; despite this, the use of seronegative ferrets is essential both for experiment reproducibility and to best assess the inherent pathogenicity and transmissibility of a given virus following primary infection. However, it must be recognised that understanding these properties in a naive animal model might not fully address the more complex real-world situation of influenza infection in human populations, which possess varying degrees of adaptive immunity to circulating strains.

Like ferrets, guinea pigs are susceptible to both human and avian influenza viruses without prior adaptation, and the utility of this model for transmission studies has recently been examined (Refs 44, 45, 46). Human influenza viruses transmit efficiently between guinea pigs by either direct contact or respiratory droplets (Refs 45, 47, 48). However, unlike in ferrets, influenza disease in guinea pigs is not clinically apparent, and thus symptoms cannot be used to assess the progression of disease among inoculated or contact animals. Furthermore, the receptor distribution of the guinea pig airway is not yet fully characterised (Refs 49, 50). Nevertheless, the relatively small size and lower cost of guinea pigs make them an attractive model for some laboratories, especially when studying the contribution of environmental conditions to virus transmissibility (Ref. 51).

Historically, mice have been the most widely used animal model for influenza virus research. Although inbred mice have been used extensively for the study of influenza virus pathogenicity, they are only infrequently used for a model of virus transmissibility. It has been hypothesised that influenza virus transmission in this model occurs primarily by airborne droplet nuclei, because comparable rates of transmission occur when infected and susceptible mice are either co-housed or separated by a barrier that allows for air exchange without direct or indirect contact between animals (Ref. 52). Transmission of mouse-adapted avian viruses in mice has also been reported in both transmission models (Ref. 53). However, a lack of consistency has been reported with this model, with a variability in results observed depending on parameters such as age of mice, virus strain tested, and route and dose of inoculum administered (Ref. 54). As such, the mouse model is not widely used to study influenza virus transmissibility. The nonhuman primate model has been used to study the virulence of select influenza viruses, but because of limited numbers, cost and ethical considerations this species is also not a routinely used model to study influenza virus transmissibility (Ref. 55).

### **Transmissibility of the 2009 H1N1 pandemic virus**

Emergence of the 2009 H1N1 virus, causing the first influenza pandemic of the 21st century,

necessitated swift study of the transmissibility, pathogenicity, and antigenic and genetic features of this strain to guide public health responses and vaccine development. 2009 H1N1 viruses were found to be antigenically similar to classical swine and North American swine-lineage H1N1 viruses but both antigenically and genetically distinct from seasonal H1N1 influenza viruses (Ref. 56). Previous studies have demonstrated the capacity of swine-origin influenza viruses to cause disease in humans; however, prior to 2009, human-to-human transmission of swine-origin H1N1 viruses was limited and not sustained (Refs 57, 58). Nevertheless, viruses of swine origin are capable of transmitting between pigs, from pigs to other mammalian species, and between mammalian species, demonstrating the ability of these viruses to pose a public health threat to humans (Refs 59, 60).

Given the efficient transmissibility in humans of seasonal influenza viruses, it was critical to determine whether 2009 H1N1 viruses shared this phenotype. Use of the ferret showed that 2009 H1N1 viruses transmitted efficiently in a direct contact model, comparable to seasonal H1N1 viruses, but with reduced efficiency by the respiratory droplet route compared with seasonal influenza viruses (Ref. 61). However, when ferrets were housed in chambers with unidirectional airflow, efficient respiratory droplet transmission was observed, demonstrating that directed airflow towards naive contacts might enhance the respiratory droplet transmissibility of this virus (Ref. 62). Decreased rates of household transmission of the 2009 H1N1 virus in the USA compared with previous pandemic viruses support the ferret modelling results, but similar rates of viral shedding, clinical illness and transmissibility between 2009 H1N1 and seasonal influenza viruses in Hong Kong suggest location-specific dynamics (Refs 63, 64). Recent studies also report low secondary transmission among household contacts of 2009-H1N1-infected individuals compared with seasonal influenza viruses (Refs 65, 66). Unlike previous pandemic viruses, 2009 H1N1 viruses do not possess known molecular determinants of virulence, and addition of these mutations to this virus strain did not confer enhanced virus replication or transmissibility (Ref. 67). Continuing surveillance of this pandemic virus will allow

for a greater understanding of how a virus capable of human-to-human transmission emerged from this constellation of avian-, human- and swine-origin genes.

### Factors affecting influenza virus transmission

Several seasonal, environmental, viral and host factors can affect the ability of influenza viruses to transmit efficiently from one host to another. Understanding the behaviour of virus-containing aerosol particles in the environment is crucial for measuring the potential of influenza virus transmission. With the advent of reverse genetics technology for influenza viruses in 1999, infectious virus could be rescued entirely from plasmid-cloned influenza gene segments without helper virus (Refs 68, 69). The reverse genetics systems used to generate reassortant viruses along with the establishment of mammalian models to study virus transmission *in vivo* have broadened our capacity to identify viral molecular determinants that confer efficient influenza virus transmission. Although reported studies are limited, there is a clear role of host genetics in the susceptibility of individuals to influenza virus infection and severe disease. These combined research efforts made it possible to rapidly assess the transmissibility of viruses isolated during the 2009 H1N1 pandemic.

### Environmental influences

Individuals infected with influenza virus generate virus-containing aerosol particles of varying sizes by a diverse range of respiratory events. Coughing, sneezing and talking are considered the principal activities that produce aerosols, but normal breathing can also generate influenza-containing aerosol particles (Ref. 70). The heterogeneity and dynamic nature of virus particles released into the environment facilitates several potential modes of exposure and subsequent transmission of virus to susceptible contacts (Table 1). Large aerosol droplets released from infected persons do not remain suspended in air and typically travel <1 m before landing on environmental surfaces or the mucosa of close contacts (Refs 27, 71). Large droplets evaporate at a slower rate than small droplets and can contain virus that might remain infectious in the environment for up to several hours, facilitating potential contact transmission of virus to a susceptible host (Refs 27, 72). In contrast, smaller particles

(<5 µm in diameter, or droplet nuclei) are capable of remaining suspended in air for longer durations of time and can be carried farther distances than large droplets, depending on the rate of particle desiccation and other environmental factors (Refs 71, 73). Particles of this size are capable of penetrating deep into the respiratory tract following inhalation, which is generally not the case for inhaled large droplets (Refs 72, 74). It is important to note that the sizes of particles described here denote generalities and do not represent absolute cutoff points for one mode of transmission or another.

Considering that size influences the movement, condensation and evaporation of aerosolised droplets generated by infected individuals, identifying the range of particle sizes expelled during respiratory events and the size range of these particles containing infectious virus is crucial for understanding the contagiousity of viral respiratory diseases (Ref. 75). For instance, variability in bioaerosol concentration and total volume in exhaled air during coughing and sneezing of children in a school setting was demonstrated by mathematical modelling to influence transmission dynamics (Ref. 76). Airway deposition, or the fraction of inhaled particles that are retained in the respiratory tract and not exhaled, varies greatly, depending on the individual's airway structure and physiology, breathing rate and volume, the size and composition of inhaled particles, and environmental conditions, such as airflow, temperature and humidity levels (Ref. 72). Once deposited in the respiratory tract, the ability of the particular virus to attach, infect and replicate at the site of deposition also has a role. Given the complexity in defining these parameters, most *in vivo* transmission studies have been limited in defining 'respiratory droplets' as inclusive of both large and small droplets in the absence of direct or indirect contact (Ref. 40). However, with increased availability of aerosol generation, sampling and analytical instrumentation, more questions addressing the behaviour of aerosols in the environment and their impact on influenza virus transmission will likely be explored (Ref. 73).

Weather conditions have been shown to further affect the persistence of aerosols in the environment and influence influenza virus transmission efficacy. It has been hypothesised that virus transmission occurs by the aerosol route in temperate climates and by the contact

**Table 1. Factors influencing behaviour of aerosols in the environment**

Factor	Effect on transmissibility	Refs
Particle size		
Distance of spread	Smaller droplets (<5 µm) travel farther distances and remain suspended in air for longer; larger droplets (>5–20 µm) generally travel <1 m and deposit on surrounding surfaces shortly after being exhaled	27, 71, 72, 73
Deposition	Smaller particles can penetrate deeper in the respiratory tract; larger particles are deposited in the upper respiratory tract	72, 74
Temperature <sup>a</sup>	Influenza viruses generally show enhanced transmissibility at cooler temperatures	72, 78, 80
Relative humidity <sup>a</sup>	Influenza viruses generally transmit with more efficiency at low relative humidity	51, 72

<sup>a</sup>As studied in experimental models.

route in tropical zones, because research using the guinea pig model has shown that aerosol but not contact transmission is sensitive to these environmental conditions (Refs 77, 78). However, only recently have studies evaluating the contribution of specific environmental conditions on influenza virus transmission been investigated in animal models (Ref. 79). Transmission studies in the guinea pig model with a seasonal H3N2 virus found that aerosol transmission efficiency was highest at 20% and 35% relative humidity, with transmission by this route abolished under conditions of high humidity (80%) (Ref. 51). This study further demonstrated efficient transmission of this H3N2 virus in guinea pigs by the aerosol route at 5°C, but not at 30°C (Ref. 51). A possible explanation for improved influenza virus transmission at lower temperatures was proposed by a group that used proton magic angle spinning nuclear magnetic resonance to generate detailed images of an influenza virus lipid envelope at different temperatures (Ref. 80). This study found that progressive ordering of lipids in the viral envelope in response to lower temperatures (like those encountered during airborne transmission in the winter months) facilitates enhanced resistance to environmental influences as it travels from one host to the next. Together, these findings point to low ambient air temperature and relative humidity as contributing factors for efficient influenza virus transmission in the winter months of temperate climates. Year-round and biannual circulation of seasonal influenza viruses in tropical and subtropical regions further underscores the importance of understanding the contribution of

environmental properties on virus transmissibility (Refs 81, 82). Further research of other parameters that are influenced by seasonality, including atmospheric–oceanic variation, host health and social behaviour such as indoor crowding during cold weather and school attendance, is needed to accurately define those conditions that contribute to the seasonality of influenza virus transmission (Refs 79, 83), especially given the unconventional global spread of the pandemic 2009 H1N1 virus during the spring and summer months.

### Contribution of receptor binding and surface glycoproteins

The influenza virus HA binds to glycoconjugates containing terminal sialic acids (SAs) present on host cells throughout the respiratory tract epithelium prior to virus entry and replication. These SA receptors are predominantly found in two linkage conformations: Neu5Acα(2,6)–Gal (abbreviated as α2–6) and Neu5Acα(2,3)–Gal (abbreviated as α2–3) (Ref. 84). Importantly, the distribution of these receptors affects the general species-specific cellular tropism of influenza viruses (Ref. 85). Human influenza viruses preferentially bind α2–6-linked SAs, which are located predominantly on nonciliated cells of the airway epithelium and nasal mucosa (Refs 61, 86, 87, 88). Avian influenza viruses, including HPAI H5N1 and H7N7 subtypes, preferentially bind α2–3-linked SAs, present on ciliated cells in the lower respiratory tract (Refs 89, 90). Studies have shown that human influenza viruses predominantly bind to upper respiratory tract epithelial cells whereas avian influenza viruses bind alveolar cells, although this differentiation


in binding is not exclusive (Refs 37, 88). Cell tropism can also be influenced by virus evolution over time; recent human H3N2 viruses show a greater tropism towards nonciliated cells compared with earlier isolates, which proportionally bind more ciliated cells (Ref. 91). Patterns of lectin binding to human airway epithelial cells demonstrate a predominance of  $\alpha$ 2–6-linked SA in the upper respiratory tract of humans (Refs 60, 61, 62). In addition to differences in linkage conformation, the structural topology of sialylated glycans further influences the binding of influenza viruses to SA (Refs 92, 93).

Molecular studies on the HA of human H1–H3 influenza viruses have revealed specific amino acids situated in the receptor-binding domain that confer a preference for binding to  $\alpha$ 2–6-linked SA. For H3 viruses, the presence of a leucine at 226 and a serine at 228 in the HA protein results in  $\alpha$ 2–6-linked SA binding; glutamine and glycine are found at these positions in avian H3 viruses and confer a binding specificity for  $\alpha$ 2–3-linked SA (Ref. 84). The receptor-binding site of the H2 HA is similar to that of H3 viruses, with a crucial role identified for residue 226 in the acquisition of human-receptor-binding specificity (Refs 94, 95). Human H1 viruses preferentially bind  $\alpha$ 2–6-linked SA but possess both Gln226 and Gly228; other HA residues, including Ser186, Asp190 and Asp225, can confer this binding preference (Refs 84, 96). The alteration of two amino acids in the HA of the reconstructed 1918 virus (Glu190 and Gly225) was sufficient to switch the H1 HA to a preference for binding  $\alpha$ 2–3-linked SA and abolish respiratory droplet transmission of this virus in the ferret model (Ref. 41) (Fig. 2). These results suggest that an important adaptation of an avian influenza virus to a mammalian host is to switch its preference for binding  $\alpha$ 2–3-linked SA.

To date, HPAI viruses of the H5 and H7 subtype have largely retained a binding specificity for  $\alpha$ 2–3-linked SA and have not demonstrated the capacity for efficient transmission through air by respiratory droplets in the ferret model (Refs 40, 43, 89, 90). Even for the rare H5 or H7 wild-type virus that possesses dual binding to  $\alpha$ 2–3- and  $\alpha$ 2–6-linked SA, there was no change in the transmission phenotype (by respiratory droplets) in ferrets (Refs 40, 89, 97). However, a North American lineage H7N2 virus with an increased affinity for  $\alpha$ 2–6-linked SA

showed efficient transmission by direct contact only in ferrets (Ref. 89). A mallard H7N3 virus able to replicate in human airway epithelial cultures *in vitro* was also capable of partial direct contact transmission in the ferret model, as was an H9N2 virus that possessed Leu226 in HA (Refs 98, 99). Although the H7 and H9 viruses discussed here did not transmit efficiently through air by respiratory droplets, the capacity for increased direct contact transmission might be the initial step towards the ultimate high-transmissible phenotype of human seasonal influenza viruses. Continued monitoring of avian influenza viruses isolated from domestic birds and humans exposed to them will provide a greater understanding of the pandemic potential of these viruses.

Site-directed mutagenesis along with reverse genetics has been used in attempts to identify the molecular changes necessary for altered the binding specificity of avian influenza viruses and to understand the effect of enhanced binding to  $\alpha$ 2–6-linked SA on avian influenza virus transmissibility. Unlike H1 viruses, substitution of Asp190 or Asp225 in the HA of an H5N1 virus is not sufficient to confer increased affinity for  $\alpha$ 2–6-linked SA (Ref. 90). Mutation of residues 226 and 228, which modulate binding specificity among H3 and H2 viruses, resulted in decreased binding of an H5N1 virus to  $\alpha$ 2–3-linked SA but only modest increases in affinity for  $\alpha$ 2–6-linked SA (Ref. 90). Substitution of additional amino acids within the HA of H5N1 viruses, including Ser193, Ala137 and Ile192, resulted in decreased binding to  $\alpha$ 2–3-linked SA or increased binding to  $\alpha$ 2–6-linked SA (Refs 100, 101). Removal of a glycosylation site (Asn158) on the H5 HA yielded similar altered binding specificity among H5N1 viruses (Ref. 101). Overall, these studies show that mutations that confer a binding preference for  $\alpha$ 2–6-linked SA among H1, H2 and H3 viruses are not sufficient to elicit comparable human binding specificity on an H5N1 virus framework. Nonetheless, the studies provide a greater understanding of the ability of H5 viruses to acquire specificity for human receptors. Whether a change in receptor-binding specificity alone is sufficient for an avian influenza virus to acquire the ability to transmit efficiently in humans is not fully known. However, in one study, recombinant H1N1 viruses were designed to test the premise that



**Figure 2. Contribution of viral molecular determinants in influenza virus transmission.** (See next page for legend.)

**Figure 2. Contribution of viral molecular determinants in influenza virus transmission.** (See previous page for figure.) Use of reverse genetics has allowed for the generation of reassortant viruses that have facilitated our understanding of which viral genes contribute towards a transmissible phenotype. Viruses were constructed, evaluated for their sialic acid binding preference and tested for their ability to transmit by respiratory droplets in the ferret model. Abbreviations: HA, haemagglutinin; M, matrix protein; NA, neuraminidase; NP, nucleoprotein; NS, nonstructural protein; PA, PB1 and PB2, polymerase proteins; RD, respiratory droplet. References: human H3N2, Refs 40, 99; human H1N1, Refs 41, 103; human H1N1 mutants, avian H1N1 mutant and avian-human H1N1 reassortants, Ref. 41; 2009 H1N1, Refs 61, 62; 2009 H1N1 mutant (Lys627), Ref. 67; H1N1 mutant (Gly222), Ref. 102; Eurasian lineage avian H5N1, Ref. 40; Eurasian lineage avian H7N7, Ref. 89; North American lineage avian H7N2, Ref. 89; Eurasian lineage avian H9N2, Ref. 99; H9N2-H2N2 reassortant, Ref. 104.

binding preference for  $\alpha$ 2-6-linked SA receptors alone would allow efficient transmission of an avian H1N1 strain. A mutant avian Dk/NY/96 H1N1 virus was constructed containing Asp190 and Asp225 (Dk/NY/96-DD) by the introduction of point mutations into the HA. Although these changes switched HA to a preference for binding  $\alpha$ 2-6-linked SA receptors, neither the Dk/NY/96-DD mutant nor the parental avian (Dk/NY/96) virus transmitted to naive ferrets in respiratory droplet experiments (Ref. 77). Conversely, the construction of a 2009 H1N1 virus containing Gly222, a substitution in the HA associated with severe human cases and fatalities, resulted in a dual binding specificity for  $\alpha$ 2-6- and  $\alpha$ 2-3-linked SA but did not alter the transmissibility of this virus compared with wild-type strains (Ref. 102).

The use of avian-human reassortant viruses has broadened our understanding of how individual genes contribute towards the overall transmissibility of influenza viruses. Substitution of genes from a transmissible human influenza virus on the framework of a nontransmissible avian influenza virus has facilitated a more detailed examination of which genes are necessary to achieve a transmissible phenotype and have shown that virus transmissibility is a complex and polygenic trait. Avian-human reassortant viruses bearing surface proteins from human virus H3N2 but with the remaining genes from an avian H5N1 virus were not capable of transmitting by respiratory droplets in the ferret model (Refs 40, 103). An H5N1 virus with internal proteins encoded by human-virus genes was similarly incapable of respiratory droplet transmission, demonstrating that the presence of human-virus surface proteins or human-virus internal proteins alone in an 'avian' virus is insufficient to achieve transmission through the air (Ref. 40). A reassortant virus bearing surface proteins of an

H9N2 virus with enhanced binding specificity for  $\alpha$ 2-6-linked SA on the framework of a human H3N2 virus was capable of direct contact transmission in ferrets; subsequent adaptation of this virus in ferrets resulted in a virus capable of transmission by respiratory droplets (Refs 99, 104). Taken together, these studies imply a crucial role for influenza virus surface glycoproteins in overall virus transmissibility, but suggest that additional virus genes are needed for efficient transmission.

#### Other viral molecular determinants affecting influenza virus transmission

Beyond HA and NA, additional viral proteins have been examined for their contributing roles in influenza virus transmission. Notably, the involvement of the influenza virus polymerase complex in virus transmissibility has been an area of active investigation. A reassortant virus bearing human influenza virus HA, NA and polymerase proteins on the framework of a nontransmissible avian influenza virus was capable of transmission by respiratory droplets in the ferret model (Ref. 103). Although reassortant viruses bearing human influenza virus surface glycoproteins and PA, PB1 or PB2 human influenza virus polymerase were all capable of efficient replication in the ferret upper respiratory tract, only the virus containing the human-virus PB2 transmitted by respiratory droplets. The PB2 protein, and residue 627 in particular, has been of particular interest given its roles in determination of host range, virulence in mammalian models and virus replication efficiency (Refs 105, 106, 107). A lysine at position 627 has been shown to enhance replication efficiency in vitro at 33°C, the temperature of the human upper respiratory tract, and viruses bearing Lys627 in PB2 replicate to higher titres in mammalian nasal passages (Refs 106, 108). The presence of Lys627

has been linked with enhanced virus transmissibility in both the ferret and guinea pig models (Refs 103, 109). In addition to Lys627, Asn701 has been associated with enhanced polymerase activity and heightened virulence, and has recently been identified as a determinant of virus transmissibility (Refs 109, 110, 111, 112).

Influenza virus infections that result in expulsion of increased amounts of respiratory fluids, by sneezing for example, might provide an increased opportunity for the transmission of virus to susceptible persons. A general correlation has been observed between viruses that predominantly bind  $\alpha$ 2–6-linked SA and the incidence of sneezing in infected ferrets (Refs 40, 103, 113). Increased replication in the upper respiratory tract could result in heightened inflammation in the epithelial lining of nasal passages and, as a result, stimulate increased mucus production and sneezing, but this has not been experimentally shown. However, efficient transmission of human influenza viruses occurs in the guinea pig model in the absence of apparent sneezing (Ref. 45). Clearly, further studies are needed, not only to identify additional molecular determinants of influenza viruses but also to better understand their influence in host responses to infection that might enhance the spread of virus among people.

### Host determinants affecting influenza virus transmission

The major focus of studies that seek to better understand influenza virus pathogenesis and transmission is on the contribution of viral determinants affecting infectivity and disease severity. Little information is available regarding host genetic determinants that might contribute to susceptibility or resistance to influenza virus infection. Nevertheless, some studies point to a potential contribution of an individual's genetic background as a contributing factor to their risk of infection and severe disease when exposed to influenza virus. Epidemiological findings have identified several family clusters of HPAI H5N1 infections, with severe outcome of infection documented among blood relatives but not among nonblood relatives such as spouses (Refs 25, 114, 115, 116). Concurrently, common hygiene practices among family members and similar exposure to contaminated environments

might also have a role (Ref. 117). A separate study investigated the risk of severe influenza infection among genealogically connected people in Utah over a 100-year period and found that close and distant relatives of individuals who died of influenza had a significantly higher risk of dying than their spousal counterparts, pointing to genetic predisposition rather than environmental similarities as the greater contributing factor (Ref. 118).

An individual's genetic makeup and immunological history undoubtedly influence the resulting immune response following virus infection and can have a strong effect on susceptibility to both infection and disease outcome. The role of immunological memory was clearly shown during the 1918 pandemic, because individuals possessing circulating HA-specific H1-virus crossreactive antibody showed enhanced survival compared with age groups likely to be serologically naive to the pandemic strain (Ref. 119). In addition to the contribution of pre-existing immunity, the interaction of viral genes with the host immune system can influence disease severity. Patients infected with the 1918 influenza virus or HPAI H5N1 viruses have been linked to a dysregulation of the inflammatory response, a result affected by both viral and host genetic determinants (Refs 120, 121, 122, 123). Additional studies that focus on genetic factors that predispose people to increased risk of infection and severe disease caused by influenza viruses are needed; recent identification of candidate genes proposed to have a role in the genetic basis for influenza virus pathogenicity will facilitate this research (Ref. 124). Because prospective human studies are usually expensive and logistically challenging, the evaluation of host genetic variations and responses to influenza virus infection has been primarily limited to the murine model (Ref. 125). A recent study demonstrated substantial changes in HPAI H5N1 virus lethality among different strains of inbred mice, identifying a role for host genetic variation in influenza-virus-induced pathology in this model (Ref. 126). As more information becomes available about genetic markers of susceptibility to influenza virus infection, we will be able to better identify at-risk groups and focus our efforts on implementing appropriate prevention and transmission control strategies.

### Pharmaceutical and nonpharmaceutical interventions and influenza virus transmission

It is evident that viral, host and environmental conditions all influence how effectively influenza A viruses spread among humans. Given this breadth of exposure variability, efforts to mitigate human-to-human transmission of influenza viruses are diverse and use both pharmaceutical and nonpharmaceutical strategies. Several therapeutic approaches to lessen disease severity and reduce virus shedding have been shown to diminish virus transmission. Although outside the scope of this review, social patterns and behaviour, including the use of face masks, hand hygiene and social distancing, can further limit virus exposure to uninfected populations. Understanding how to maximise the effectiveness of each approach will contribute towards decreased influenza virus infection rates during both pandemic and inter-pandemic times.

### Vaccination

Vaccination is the primary public health strategy to prevent or reduce disease following influenza virus infection. Simulation models demonstrate that vaccination has the potential to significantly reduce the spread and severity of disease in the event of a pandemic, although these transmission dynamics might be influenced by prior host immunity to heterologous strains (Refs 127, 128). Two vaccine formulations are licensed for use in the USA: an inactivated virus preparation and a live attenuated virus preparation (Refs 129, 130). Clinical trials have shown reduced incidence of influenza infection among contacts of vaccinated individuals, suggesting that vaccination interrupts the transmission chain by bolstering herd immunity (Refs 131, 132). Vaccination of healthcare workers in long-term care facilities has also been associated with decreased rates of virus infection and mortality among patients (Refs 133, 134). However, the ability of vaccines to specifically reduce virus transmission has not been comprehensively examined. A recent study in the guinea pig model demonstrated that influenza virus vaccination does not uniformly block virus transmission following challenge (Ref. 135). Intranasally administered live attenuated vaccine elicited immunity in guinea pigs against both homologous and heterologous H3N2 viruses, effectively preventing virus

transmission to naive contacts. Although guinea pigs vaccinated intramuscularly with inactivated vaccine showed reduced viral loads compared with unvaccinated animals, a complete block of virus transmission to contact animals was not observed (Ref. 135). These results indicate that the ability of a vaccine to protect against virus infection does not necessarily correlate with the ability to block virus transmission following vaccination. More extensive evaluation of vaccine candidates to examine their ability to mitigate both disease symptoms and transmission would offer a more complete understanding of vaccine efficacy.

### Antiviral drugs

The use of antiviral drugs is considered a first line of defence in the event of a pandemic, and simulation modelling has demonstrated the potential of targeted prophylaxis antiviral use to contain the spread of a novel influenza virus (Ref. 136). There are currently two classes of antiviral drugs licensed for the treatment of influenza: adamantanes and NA inhibitors. The drug amantadine and its analogue rimantadine function by blocking the ion channel formed by the viral M2 protein, thus altering endosomal pH levels and inhibiting the HA-mediated fusion of viral and endosomal membranes (Refs 137, 138). However, viral resistance to this class of drugs can be achieved by acquisition of key mutations in the drug target site, the transmembrane region of the M2 protein (Ref. 139). Studies in the ferret model have demonstrated that seasonal influenza viruses that acquire adamantane resistance maintain the virulence and replication efficiency of wild-type virus without a loss of fitness (Refs 140, 141). Drug-resistant strains appear to transmit as readily as sensitive strains, and the high prevalence of resistant viruses in circulation has unfortunately limited the use of this class of antiviral drugs (Refs 142, 143). Two inhibitors of viral NA – oseltamivir and zanamivir – are effective against both human and avian influenza viruses because neutralisation of sialidase activity results in the aggregation of virus particles to SA on the surface of infected cells, preventing the effective release of virions and spread to uninfected cells (Refs 144, 145). Resistance to this class of antiviral drugs occurs less frequently compared with the adamantanes, because mutations in the NA active site are both

largely subtype specific and more likely to adversely affect the function of this protein. Several key mutations in NA that confer a drug-resistant phenotype, including Val119 (H3N2), Lys292 (H3N2) and Tyr274 (H1N1), show decreased viral fitness in the ferret model, with reduced infectivity, replication and pathogenicity of seasonal influenza viruses compared with viruses that do not bear these changes (Refs 146, 147, 148). HPAI H5N1 viruses, which do not readily transmit in the ferret model, are capable of maintaining their high pathogenicity phenotype on the introduction of mutations known to confer NA resistance (Ref. 149).

Although mutations in NA frequently compromise overall viral fitness, the transmissibility of viruses resistant to NA inhibitors is dependent on the specific mutations conferring a resistant phenotype. Viruses with Val119 and Tyr274 mutations in NA are capable of direct contact transmission in the ferret model, unlike viruses that possess Lys292 in this protein (Refs 148, 150). However, viruses possessing the Tyr274 mutation require higher challenge doses to transmit, and studies in the guinea pig model have shown that Val119 viruses show decreased transmission by the aerosol route compared with wild-type viruses (Ref. 151). This observed variation in virus fitness and transmissibility in the ferret model appears to be dependent on the degree of NA functional loss (Ref. 152). Study of these mutations on a homogeneous genetic background showed that NA mutations conferring greater reductions of enzymatic activity and thermostability, such as Lys292, show greater reductions in overall transmissibility compared with viruses that retain these properties, such as viruses possessing Val119 (Ref. 152). This work indicates that viruses with mutations facilitating resistance to NA drugs are capable of being transmitted among people, albeit at reduced efficiency compared with wild-type viruses that retain sensitivity to these antivirals. The stockpiling of antiviral drugs is used by several nations (Ref. 153); in addition to concerns of quantity, drug resistance and availability of paediatric formulations, the effectiveness of these formulations in reducing the transmission of new virus strains among treated individuals should be evaluated and there needs to be continued surveillance to identify any influenza virus strains that acquire resistance to available antivirals.

Postexposure oseltamivir prophylaxis of contacts of individuals with laboratory-confirmed influenza reduced the secondary spread of virus to susceptible individuals in two recent studies, demonstrating the ability of antiviral treatment to attenuate virus transmission (Refs 154, 155). Furthermore, additional antiviral drugs need to be developed in the event that influenza viruses resistant to NA inhibitors attain increased circulation and pose a public health risk.

### Interferon treatment in preclinical models

The host interferon (IFN) response has an important role in the establishment of an antiviral state that can limit virus replication following infection. Administration of exogenous IFN to engage this response prior to infection is a potential prophylactic or therapeutic strategy for influenza viruses and is used clinically for the treatment of other viral infections, including hepatitis C virus (Refs 156, 157). Previous studies have demonstrated the ability of exogenous IFN to reduce viral titres and to reduce disease severity following influenza virus infection in several animal models, including mice, ferrets and guinea pigs (Refs 158, 159, 160). It was recently shown in the guinea pig model that intranasal treatment with human IFN- $\alpha$  initiated one day prior to viral infection significantly reduced or prevented infection with a highly pathogenic H5N1 virus, the reconstructed 1918 pandemic virus, or the 2009 pandemic virus (Refs 48, 161). Furthermore, treatment of either inoculated or contact guinea pigs with IFN was sufficient to prevent transmission of the 2009 pandemic virus by direct contact (Ref. 161). This work indicates that IFN treatment is capable of reducing viral titres and disease symptoms in infected animals in addition to blocking the transmission of virus to naive animals. These results suggest that investigations are warranted to assess the potential of IFN treatment for clinical use.

### Conclusions

Despite these and other research efforts, much remains in question regarding the viral and host determinants of efficient influenza virus transmission. The seasonality of influenza virus transmission highlights the importance of environmental conditions on transmission efficiency, and with increased availability of specialised instrumentation to test the

environmental parameters affecting influenza-virus-containing aerosols, many of these unanswered questions can be addressed (Ref. 79). Understanding the mechanisms involved in influenza virus infection, transmission and disease progression will facilitate the implementation of more-effective clinical prevention and treatment strategies, preparing us not only for the next influenza season but also for the next influenza pandemic.

### Acknowledgements and funding

We thank the peer reviewers of this manuscript for helpful suggestions. The authors have no financial conflicts to report. The findings and conclusions in this report are those of the authors and do not necessarily reflect the views of the funding agency.

### References

- 1 Chen, W. et al. (2001) A novel influenza A virus mitochondrial protein that induces cell death. *Nature Medicine* 7, 1306-1312
- 2 Webster, R.G. et al. (1992) Evolution and ecology of influenza A viruses. *Microbiology Reviews* 56, 152-179
- 3 Fouchier, R.A. et al. (2005) Characterization of a novel influenza A virus hemagglutinin subtype (H16) obtained from black-headed gulls. *Journal of Virology* 79, 2814-2822
- 4 Abdel-Ghafar, A.N. et al. (2008) Update on avian influenza A (H5N1) virus infection in humans. *New England Journal of Medicine* 358, 261-273
- 5 Belser, J.A. et al. (2009) Past, present, and possible future human infection with influenza virus A subtype H7. *Emerging Infectious Diseases* 15, 859-865
- 6 Alford, R.H. et al. (1966) Human influenza resulting from aerosol inhalation. *Proceedings of the Society for Experimental Biology and Medicine* 122, 800-804
- 7 Bean, B. et al. (1982) Survival of influenza viruses on environmental surfaces. *Journal of Infectious Diseases* 146, 47-51
- 8 Lidwell, O.M. (1974) Aerial dispersal of micro-organisms from the human respiratory tract. *Society for Applied Bacteriology Symposium Series* 3, 135-154
- 9 Potter, W.P. (1998) Chronicle of influenza pandemics. In *Textbook of Influenza* (Nicholson K.G., Webster R.G. and Hay A.J., eds), pp. 3-18, Blackwell Science, Oxford
- 10 Miller, J.M. et al. (2000) Cruise ships: high-risk passengers and the global spread of new influenza viruses. *Clinical Infectious Diseases* 31, 433-438
- 11 Mizuta, K. et al. (1995) An outbreak of influenza A/H3N2 in a Zambian school dormitory. *East African Medical Journal* 72, 189-190
- 12 Earhart, K.C. et al. (2001) Outbreak of influenza in highly vaccinated crew of U.S. Navy ship. *Emerging Infectious Diseases* 7, 463-465
- 13 Horcajada, J.P. et al. (2003) A nosocomial outbreak of influenza during a period without influenza epidemic activity. *European Respiratory Journal* 21, 303-307
- 14 Horman, J.T. et al. (1986) An outbreak of influenza A in a nursing home. *American Journal of Public Health* 76, 501-504
- 15 Moser, M.R. et al. (1979) An outbreak of influenza aboard a commercial airliner. *American Journal of Epidemiology* 110, 1-6
- 16 Leder, K. and Newman, D. (2005) Respiratory infections during air travel. *Internal Medicine Journal* 35, 50-55
- 17 Khan, K. et al. (2009) Spread of a novel influenza A (H1N1) virus via global airline transportation. *New England Journal of Medicine* 361, 212-214
- 18 Taubenberger, J.K. and Morens, D.M. (2006) 1918 Influenza: the mother of all pandemics. *Emerging Infectious Diseases* 12, 15-22
- 19 Thompson, W.W. et al. (2004) Influenza-associated hospitalizations in the United States. *Journal of the American Medical Association* 292, 1333-1340
- 20 Beare, A.S. and Webster, R.G. (1991) Replication of avian influenza viruses in humans. *Archives of Virology* 119, 37-42
- 21 Uyeki, T.M. et al. (2002) Lack of evidence for human-to-human transmission of avian influenza A (H9N2) viruses in Hong Kong, China 1999. *Emerging Infectious Diseases* 8, 154-159
- 22 Peiris, M. et al. (1999) Human infection with influenza H9N2. *Lancet* 354, 916-917
- 23 Koopmans, M. et al. (2004) Transmission of H7N7 avian influenza A virus to human beings during a large outbreak in commercial poultry farms in the Netherlands. *Lancet* 363, 587-593
- 24 Kandun, I.N. et al. (2006) Three Indonesian clusters of H5N1 virus infection in 2005. *New England Journal of Medicine* 355, 2186-2194
- 25 Olsen, S.J. et al. (2005) Family clustering of avian influenza A (H5N1). *Emerging Infectious Diseases* 11, 1799-1801
- 26 Ungchusak, K. et al. (2005) Probable person-to-person transmission of avian influenza A (H5N1). *New England Journal of Medicine* 352, 333-340
- 27 Brankston, G. et al. (2007) Transmission of influenza A in human beings. *Lancet Infectious Diseases* 7, 257-265

- 28 Mubareka, S. et al. (2009) Transmission of influenza virus via aerosols and fomites in the guinea pig model. *Journal of Infectious Diseases* 199, 858-865
- 29 Uyeki, T.M. (2009) Human infection with highly pathogenic avian influenza A (H5N1) virus: review of clinical issues. *Clinical Infectious Diseases* 49, 279-290
- 30 Rimmelzwaan, G.F. et al. (2006) Influenza A virus (H5N1) infection in cats causes systemic disease with potential novel routes of virus spread within and between hosts. *American Journal of Pathology* 168, 176-183; quiz 364
- 31 Lipatov, A.S. et al. (2009) Pathogenesis of H5N1 influenza virus infections in mice and ferret models differs according to respiratory tract or digestive system exposure. *Journal of Infectious Diseases* 199, 717-725
- 32 Olofsson, S. et al. (2005) Avian influenza and sialic acid receptors: more than meets the eye? *Lancet Infectious Diseases* 5, 184-188
- 33 Du Ry van Beest Holle, M. et al. (2005) Human-to-human transmission of avian influenza A/H7N7, The Netherlands, 2003. *Euro Surveillance* 10, 264-268
- 34 Smith, H. and Sweet, C. (1988) Lessons for human influenza from pathogenicity studies with ferrets. *Review of Infectious Diseases* 10, 56-75
- 35 Sweet, C. et al. (1979) The local origin of the febrile response induced in ferrets during respiratory infection with a virulent influenza virus. *British Journal of Experimental Pathology* 60, 300-308
- 36 van Riel, D. et al. (2007) Human and avian influenza viruses target different cells in the lower respiratory tract of humans and other mammals. *American Journal of Pathology* 171, 1215-1223
- 37 van Riel, D. et al. (2006) H5N1 Virus Attachment to Lower Respiratory Tract. *Science* 312, 399
- 38 Andrewes, C.H. and Glover, R.E. (1941) Spread of infection from the respiratory tract of the ferret. I. Transmission of influenza A virus. *British Journal of Experimental Pathology* 22, 91
- 39 Herlocher, M.L. et al. (2001) Ferrets as a transmission model for influenza: sequence changes in HA1 of type A (H3N2) virus. *Journal of Infectious Diseases* 184, 542-546
- 40 Maines, T.R. et al. (2006) Lack of transmission of H5N1 avian-human reassortant influenza viruses in a ferret model. *Proceedings of the National Academy of Sciences of the United States of America* 103, 12121-12126
- 41 Tumpey, T.M. et al. (2007) A two-amino acid change in the hemagglutinin of the 1918 influenza virus abolishes transmission. *Science* 315, 655-659
- 42 Squires, S. and Belyavin, G. (1975) Free contact infection in ferret groups. *Journal of Antimicrobial Chemotherapy* 1 (Supplement 4), 35-42
- 43 Yen, H.L. et al. (2007) Inefficient transmission of H5N1 influenza viruses in a ferret contact model. *Journal of Virology* 81, 6890-6898
- 44 Azoulay-Dupuis, E. et al. (1984) Lung alterations in guinea-pigs infected with influenza virus. *Journal of Comparative Pathology* 94, 273-283
- 45 Lowen, A.C. et al. (2006) The guinea pig as a transmission model for human influenza viruses. *Proceedings of the National Academy of Sciences of the United States of America* 103, 9988-9992
- 46 Wetherbee, R.E. (1973) Induction of systemic delayed hypersensitivity during experimental viral infection of the respiratory tract with a myxovirus or paramyxovirus. *Journal of Immunology* 111, 157-163
- 47 Lowen, A.C. and Palese, P. (2007) Influenza virus transmission: basic science and implications for the use of antiviral drugs during a pandemic. *Infectious Disorders Drug Targets* 7, 318-328
- 48 Van Hoeven, N. et al. (2009) Pathogenesis of 1918 pandemic and H5N1 influenza virus infections in a guinea pig model: antiviral potential of exogenous alpha interferon to reduce virus shedding. *Journal of Virology* 83, 2851-2861
- 49 Pastor, L.M. et al. (1992) Histochemical study of glycoconjugates in the nasal mucosa of the rat and guinea pig. *Histochemical Journal* 24, 727-736
- 50 Jarreau, P.H. et al. (1992) Effects of neuraminidase on airway reactivity in the guinea pig. *American Review of Respiratory Disease* 145(4 Pt 1), 906-910
- 51 Lowen, A.C. et al. (2007) Influenza virus transmission is dependent on relative humidity and temperature. *PLoS Pathogens* 3, 1470-1476
- 52 Schulman, J.L. and Kilbourne, E.D. (1962) Airborne transmission of influenza virus infection in mice. *Nature* 195, 1129-1130
- 53 Wu, R. et al. (2010) Transmission of avian H9N2 influenza viruses in a murine model. *Veterinary Microbiology* 142, 211-216
- 54 Schulman, J.L. and Kilbourne, E.D. (1963) Experimental transmission of influenza virus infection in mice. II. Some factors affecting the incidence of transmitted infection. *Journal of Experimental Medicine* 118, 267-275
- 55 Baskin, C.R. et al. (2009) Early and sustained innate immune response defines pathology and death in nonhuman primates infected by highly pathogenic influenza virus. *Proceedings of the National Academy of Sciences of the United States of America* 106, 3455-3460

- 56 Garten, R.J. et al. (2009) Antigenic and genetic characteristics of swine-origin 2009 A(H1N1) influenza viruses circulating in humans. *Science* 325, 197-201
- 57 Myers, K.P., Olsen, C.W. and Gray, G.C. (2007) Cases of swine influenza in humans: a review of the literature. *Clinical Infectious Diseases* 44, 1084-1088
- 58 Shinde, V. et al. (2009) Triple-reassortant swine influenza A (H1) in humans in the United States, 2005-2009. *New England Journal of Medicine* 360, 2616-2625
- 59 De Vleeschauwer, A. et al. (2009) Efficient transmission of swine-adapted but not wholly avian influenza viruses among pigs and from pigs to ferrets. *Journal of Infectious Diseases* 200, 1884-1892
- 60 Ma, W. et al. (2007) Identification of H2N3 influenza A viruses from swine in the United States. *Proceedings of the National Academy of Sciences of the United States of America* 104, 20949-20954
- 61 Maines, T.R. et al. (2009) Transmission and pathogenesis of swine-origin 2009 A(H1N1) influenza viruses in ferrets and mice. *Science* 325, 484-487
- 62 Munster, V.J. et al. (2009) Pathogenesis and transmission of swine-origin 2009 A(H1N1) influenza virus in ferrets. *Science* 325, 481-483
- 63 Cauchemez, S. et al. (2009) Household transmission of 2009 pandemic influenza A (H1N1) virus in the United States. *New England Journal of Medicine* 361, 2619-2627
- 64 Cowling, B.J. et al. (2010) Comparative epidemiology of pandemic and seasonal influenza A in households. *New England Journal of Medicine* 362, 2175-2184
- 65 Doyle, T.J. and Hopkins, R.S. (2010) Low secondary transmission of 2009 pandemic influenza A (H1N1) in households following an outbreak at a summer camp: relationship to timing of exposure. *Epidemiology and Infection* Jun 21; [Epub ahead of print]
- 66 Leung, Y.H., Li, M.P. and Chuang, S.K. (2010) A school outbreak of pandemic (H1N1) 2009 infection: assessment of secondary household transmission and the protective role of oseltamivir. *Epidemiology and Infection* Jun 21; [Epub ahead of print]
- 67 Herfst, S. et al. (2010) Introduction of virulence markers in PB2 of pandemic swine-origin influenza virus does not result in enhanced virulence or transmission. *Journal of Virology* 84, 3752-3758
- 68 Hoffmann, E. et al. (2000) A DNA transfection system for generation of influenza A virus from eight plasmids. *Proceedings of the National Academy of Sciences of the United States of America* 97, 6108-6113
- 69 Fodor, E. et al. (1999) Rescue of influenza A virus from recombinant DNA. *Journal of Virology* 73, 9679-9682
- 70 Stelzer-Braid, S. et al. (2009) Exhalation of respiratory viruses by breathing, coughing, and talking. *Journal of Medical Virology* 81, 1674-1679
- 71 Nicas, M., Nazaroff, W.W. and Hubbard, A. (2005) Toward understanding the risk of secondary airborne infection: emission of respirable pathogens. *Journal of Occupational and Environmental Hygiene* 2, 143-154
- 72 Hinds, W.C. (1999) *Aerosol Technology* (2nd edn), John Wiley & Sons, Inc., New York
- 73 Verreault, D., Moineau, S. and Duchaine, C. (2008) Methods for sampling of airborne viruses. *Microbiology and Molecular Biology Reviews* 72, 413-444
- 74 Tellier, R. (2009) Aerosol transmission of influenza A virus: a review of new studies. *Journal of the Royal Society, Interface* 6 (Supplement 6), S783-S790
- 75 Xie, X. et al. (2007) How far droplets can move in indoor environments – revisiting the Wells evaporation-falling curve. *Indoor Air* 17, 211-225
- 76 Chen, S.C. et al. (2009) Viral kinetics and exhaled droplet size affect indoor transmission dynamics of influenza infection. *Indoor Air* 19, 401-413
- 77 Lowen, A. and Palese, P. (2009) Transmission of influenza virus in temperate zones is predominantly by aerosol, in the tropics by contact: a hypothesis. *PLoS Currents: Influenza* Aug 17; 1:RRN1002
- 78 Lowen, A.C. et al. (2008) High temperature (30 degrees C) blocks aerosol but not contact transmission of influenza virus. *Journal of Virology* 82, 5650-5652
- 79 Lofgren, E. et al. (2007) Influenza seasonality: underlying causes and modeling theories. *Journal of Virology* 81, 5429-5436
- 80 Polozov, I.V. et al. (2008) Progressive ordering with decreasing temperature of the phospholipids of influenza virus. *Nature Chemical Biology* 4, 248-255
- 81 Nguyen, H.L. et al. (2007) Epidemiology of influenza in Hanoi, Vietnam, from 2001 to 2003. *Journal of Infection* 55, 58-63
- 82 Wong, C.M. et al. (2004) Influenza-associated mortality in Hong Kong. *Clinical Infectious Diseases* 39, 1611-1617
- 83 Viboud, C. et al. (2004) Association of influenza epidemics with global climate variability. *European Journal of Epidemiology* 19, 1055-1059

- 84 Skehel, J.J. and Wiley, D.C. (2000) Receptor binding and membrane fusion in virus entry: the influenza hemagglutinin. *Annual Review of Biochemistry* 69, 531-569
- 85 Ito, T. and Kawaoka, Y. (2000) Host-range barrier of influenza A viruses. *Veterinary Microbiology* 74, 71-75
- 86 Rogers, G.N. and Paulson, J.C. (1983) Receptor determinants of human and animal influenza virus isolates: differences in receptor specificity of the H3 hemagglutinin based on species of origin. *Virology* 127, 361-373
- 87 Matrosovich, M.N. et al. (2004) Human and avian influenza viruses target different cell types in cultures of human airway epithelium. *Proceedings of the National Academy of Sciences of the United States of America* 101, 4620-4624
- 88 Shinya, K. et al. (2006) Avian flu: influenza virus receptors in the human airway. *Nature* 440, 435-436
- 89 Belser, J.A. et al. (2008) Contemporary North American influenza H7 viruses possess human receptor specificity: implications for virus transmissibility. *Proceedings of the National Academy of Sciences of the United States of America* 105, 7558-7563
- 90 Stevens, J. et al. (2006) Structure and receptor specificity of the hemagglutinin from an H5N1 influenza virus. *Science* 312, 404-410
- 91 Thompson, C.I. et al. (2006) Infection of human airway epithelium by human and avian strains of influenza A virus. *Journal of Virology* 80, 8060-8068
- 92 Chandrasekaran, A. et al. (2008) Glycan topology determines human adaptation of avian H5N1 virus hemagglutinin. *Nature Biotechnology* 26, 107-113
- 93 Srinivasan, A. et al. (2008) Quantitative biochemical rationale for differences in transmissibility of 1918 pandemic influenza A viruses. *Proceedings of the National Academy of Sciences of the United States of America* 105, 2800-2805
- 94 Xu, R. et al. (2010) Structure, receptor binding, and antigenicity of influenza virus hemagglutinins from the 1957 H2N2 pandemic. *Journal of Virology* 84, 1715-1721
- 95 Liu, J. et al. (2009) Structures of receptor complexes formed by hemagglutinins from the Asian influenza pandemic of 1957. *Proceedings of the National Academy of Sciences of the United States of America* 106, 17175-17180
- 96 Glaser, L. et al. (2005) A single amino acid substitution in 1918 influenza virus hemagglutinin changes receptor binding specificity. *Journal of Virology* 79, 11533-11536
- 97 Gambaryan, A. et al. (2006) Evolution of the receptor binding phenotype of influenza A (H5) viruses. *Virology* 344, 432-438
- 98 Song, H. et al. (2009) Partial direct contact transmission in ferrets of a mallard H7N3 influenza virus with typical avian-like receptor specificity. *Virology Journal* 6, 126
- 99 Wan, H. et al. (2008) Replication and transmission of H9N2 influenza viruses in ferrets: evaluation of pandemic potential. *PLoS One* 3, e2923
- 100 Yang, Z.Y. et al. (2007) Immunization by avian H5 influenza hemagglutinin mutants with altered receptor binding specificity. *Science* 317, 825-828
- 101 Stevens, J. et al. (2008) Recent avian H5N1 viruses exhibit increased propensity for acquiring human receptor specificity. *Journal of Molecular Biology* 381, 1382-1394
- 102 Chutinimitkul, S. et al. (2010) Virulence-associated substitution D222G in hemagglutinin of 2009 pandemic influenza A(H1N1) virus affects receptor binding. *Journal of Virology* 84, 11802-11813
- 103 Van Hoeven, N. et al. (2009) Human HA and polymerase subunit PB2 proteins confer transmission of an avian influenza virus through the air. *Proceedings of the National Academy of Sciences of the United States of America* 106, 3366-3371
- 104 Sorrell, E.M. et al. (2009) Minimal molecular constraints for respiratory droplet transmission of an avian-human H9N2 influenza A virus. *Proceedings of the National Academy of Sciences of the United States of America* 106, 7565-7570
- 105 Subbarao, E.K., London, W. and Murphy, B.R. (1993) A single amino acid in the PB2 gene of influenza A virus is a determinant of host range. *Journal of Virology* 67, 1761-1764
- 106 Massin, P., van der Werf, S. and Naffakh, N. (2001) Residue 627 of PB2 is a determinant of cold sensitivity in RNA replication of avian influenza viruses. *Journal of Virology* 75, 5398-5404
- 107 Hatta, M. et al. (2001) Molecular basis for high virulence of Hong Kong H5N1 influenza A viruses. *Science* 293, 1840-1842
- 108 Hatta, M. et al. (2007) Growth of H5N1 influenza A viruses in the upper respiratory tracts of mice. *PLoS Pathogens* 3, 1374-1379
- 109 Steel, J. et al. (2009) Transmission of influenza virus in a mammalian host is increased by PB2 amino acids 627K or 627E/701N. *PLoS Pathogens* 5, e1000252
- 110 Li, Z. et al. (2005) Molecular basis of replication of duck H5N1 influenza viruses in a mammalian mouse model. *Journal of Virology* 79, 12058-12064

- 111 Gabriel, G. et al. (2005) The viral polymerase mediates adaptation of an avian influenza virus to a mammalian host. *Proceedings of the National Academy of Sciences of the United States of America* 102, 18590-18595
- 112 Gao, Y. et al. (2009) Identification of amino acids in HA and PB2 critical for the transmission of H5N1 avian influenza viruses in a mammalian host. *PLoS Pathogens* 5, e1000709
- 113 Leigh, M.W. et al. (1995) Receptor specificity of influenza virus influences severity of illness in ferrets. *Vaccine* 13, 1468-1473
- 114 Dinh, P.N. et al. (2006) Risk factors for human infection with avian influenza A H5N1, Vietnam, 2004. *Emerging Infectious Diseases* 12, 1841-1847
- 115 Normile, D. (2007) Epidemiology. Indonesia taps village wisdom to fight bird flu. *Science* 315, 30-33
- 116 Beigel, J.H. et al. (2005) Avian influenza A (H5N1) infection in humans. *New England Journal of Medicine* 353, 1374-1385
- 117 Pitzer, V.E. et al. (2007) Little evidence for genetic susceptibility to influenza A (H5N1) from family clustering data. *Emerging Infectious Diseases* 13, 1074-1076
- 118 Albright, F.S. et al. (2008) Evidence for a heritable predisposition to death due to influenza. *Journal of Infectious Diseases* 197, 18-24
- 119 Ahmed, R., Oldstone, M.B. and Palese, P. (2007) Protective immunity and susceptibility to infectious diseases: lessons from the 1918 influenza pandemic. *Nature Immunology* 8, 1188-1193
- 120 Yuen, K.Y. et al. (1998) Clinical features and rapid viral diagnosis of human disease associated with avian influenza A H5N1 virus. *Lancet* 351, 467-471
- 121 Peiris, J.S. et al. (2004) Re-emergence of fatal human influenza A subtype H5N1 disease. *Lancet* 363, 617-619
- 122 Cheung, C.Y. et al. (2002) Induction of proinflammatory cytokines in human macrophages by influenza A (H5N1) viruses: a mechanism for the unusual severity of human disease? *Lancet* 360, 1831-1837
- 123 de Jong, M.D. et al. (2006) Fatal outcome of human influenza A (H5N1) is associated with high viral load and hypercytokinemia. *Nature Medicine* 12, 1203-1207
- 124 Zhang, L. et al. (2009) Systems-based candidate genes for human response to influenza infection. *Infection, Genetics and Evolution* 9, 1148-1157
- 125 Trammell, R.A. and Toth, L.A. (2008) Genetic susceptibility and resistance to influenza infection and disease in humans and mice. *Expert Review of Molecular Diagnostics* 8, 515-529
- 126 Boon, A.C. et al. (2009) Host genetic variation affects resistance to infection with a highly pathogenic H5N1 influenza A virus in mice. *Journal of Virology* 83, 10417-10426
- 127 Germann, T.C. et al. (2006) Mitigation strategies for pandemic influenza in the United States. *Proceedings of the National Academy of Sciences of the United States of America* 103, 5935-5940
- 128 Park, A.W. et al. (2009) Quantifying the impact of immune escape on transmission dynamics of influenza. *Science* 326, 726-728
- 129 Bridges, C.B. et al. (2008) Inactivated influenza vaccines. In *Vaccines (5th edn)* (Plotkin S.A., Orenstein W.A. and Offit P.A., eds), pp. 259-290, Elsevier
- 130 Belshe, R.B. et al. (2008) Influenza vaccine-live. In *Vaccines (5th edn)* (Plotkin S.A., Orenstein W.A. and Offit P.A., eds), pp. 291-310, Elsevier
- 131 Piedra, P.A. et al. (2005) Herd immunity in adults against influenza-related illnesses with use of the trivalent-live attenuated influenza vaccine (CAIV-T) in children. *Vaccine* 23, 1540-1548
- 132 Glezen, W.P. (2006) Herd protection against influenza. *Journal of Clinical Virology* 37, 237-243
- 133 Potter, J. et al. (1997) Influenza vaccination of health care workers in long-term-care hospitals reduces the mortality of elderly patients. *Journal of Infectious Diseases* 175, 1-6
- 134 Carman, W.F. et al. (2000) Effects of influenza vaccination of health-care workers on mortality of elderly people in long-term care: a randomised controlled trial. *Lancet* 355, 93-97
- 135 Lowen, A.C. et al. (2009) Blocking interhost transmission of influenza virus by vaccination in the guinea pig model. *Journal of Virology* 83, 2803-2818
- 136 Longini, I.M. Jr., et al. (2004) Containing pandemic influenza with antiviral agents. *American Journal of Epidemiology* 159, 623-633
- 137 Pinto, L.H., Holsinger, L.J. and Lamb, R.A. (1992) Influenza virus M2 protein has ion channel activity. *Cell* 69, 517-528
- 138 Hay, A.J. et al. (1985) The molecular basis of the specific anti-influenza action of amantadine. *EMBO Journal* 4, 3021-3024
- 139 Simonsen, L. et al. (2007) The genesis and spread of reassortment human influenza A/H3N2 viruses conferring adamantane resistance. *Molecular Biology and Evolution* 24, 1811-1820
- 140 Herlocher, M.L. et al. (2003) Assessment of development of resistance to antivirals in the ferret model of influenza virus infection. *Journal of Infectious Diseases* 188, 1355-1361

- 141 Sweet, C. et al. (1991) Virulence of rimantadine-resistant human influenza A (H3N2) viruses in ferrets. *Journal of Infectious Diseases* 164, 969-972
- 142 Hayden, F.G. et al. (1989) Emergence and apparent transmission of rimantadine-resistant influenza A virus in families. *New England Journal of Medicine* 321, 1696-1702
- 143 Bright, R.A. et al. (2006) Adamantane resistance among influenza A viruses isolated early during the 2005-2006 influenza season in the United States. *Journal of the American Medical Association* 295, 891-894
- 144 von Itzstein, M. et al. (1993) Rational design of potent sialidase-based inhibitors of influenza virus replication. *Nature* 363, 418-423
- 145 Gubareva, L.V., Kaiser, L. and Hayden, F.G. (2000) Influenza virus neuraminidase inhibitors. *Lancet* 355, 827-835
- 146 Carr, J. et al. (2002) Influenza virus carrying neuraminidase with reduced sensitivity to oseltamivir carboxylate has altered properties in vitro and is compromised for infectivity and replicative ability in vivo. *Antiviral Research* 54, 79-88
- 147 Ives, J.A. et al. (2002) The H274Y mutation in the influenza A/H1N1 neuraminidase active site following oseltamivir phosphate treatment leave virus severely compromised both in vitro and in vivo. *Antiviral Research* 55, 307-317
- 148 Herlocher, M.L. et al. (2002) Influenza virus carrying an R292K mutation in the neuraminidase gene is not transmitted in ferrets. *Antiviral Research* 54, 99-111
- 149 Yen, H.L. et al. (2007) Neuraminidase inhibitor-resistant recombinant A/Vietnam/1203/04 (H5N1) influenza viruses retain their replication efficiency and pathogenicity in vitro and in vivo. *Journal of Virology* 81, 12418-12426
- 150 Herlocher, M.L. et al. (2004) Influenza viruses resistant to the antiviral drug oseltamivir: transmission studies in ferrets. *Journal of Infectious Diseases* 190, 1627-1630
- 151 Bouvier, N.M., Lowen, A.C. and Palese, P. (2008) Oseltamivir-resistant influenza A viruses are transmitted efficiently among guinea pigs by direct contact but not by aerosol. *Journal of Virology* 82, 10052-10058
- 152 Yen, H.L. et al. (2005) Neuraminidase inhibitor-resistant influenza viruses may differ substantially in fitness and transmissibility. *Antimicrobial Agents and Chemotherapy* 49, 4075-4084
- 153 Patel, A. and Gorman, S.E. (2009) Stockpiling antiviral drugs for the next influenza pandemic. *Clinical Pharmacology and Therapeutics* 86, 241-243
- 154 Hayden, F.G. et al. (2004) Management of influenza in households: a prospective, randomized comparison of oseltamivir treatment with or without postexposure prophylaxis. *Journal of Infectious Diseases* 189, 440-449
- 155 Lee, V.J. et al. (2010) Oseltamivir ring prophylaxis for containment of 2009 H1N1 influenza outbreaks. *New England Journal of Medicine* 362, 2166-2174
- 156 Garcia-Sastre, A. and Biron, C.A. (2006) Type 1 interferons and the virus-host relationship: a lesson in detente. *Science* 312, 879-882
- 157 Keam, S.J. and Cvetkovic, R.S. (2008) Peginterferon-alpha-2a (40 kD) plus ribavirin: a review of its use in the management of chronic hepatitis C mono-infection. *Drugs* 68, 1273-1317
- 158 Beilharz, M.W., Cummins, J.M. and Bennett, A.L. (2007) Protection from lethal influenza virus challenge by oral type 1 interferon. *Biochemical and Biophysical Research Communications* 355, 740-744
- 159 Kugel, D. et al. (2009) Intranasal administration of alpha interferon reduces seasonal influenza A virus morbidity in ferrets. *Journal of Virology* 83, 3843-3851
- 160 Tumpey, T.M. et al. (2007) The Mx1 gene protects mice against the pandemic 1918 and highly lethal human H5N1 influenza viruses. *Journal of Virology* 81, 10818-10821
- 161 Steel, J. et al. (2010) Transmission of pandemic H1N1 influenza virus and impact of prior exposure to seasonal strains or interferon treatment. *Journal of Virology* 84, 21-26

### Further reading, resources and contacts

Tellier, R. (2009) Aerosol transmission of influenza A virus: a review of new studies. *Journal of the Royal Society, Interface* 6 (Suppl. 6), S783-S790  
A recent review of laboratory, clinical and epidemiological studies assessing the role of influenza virus transmission by the aerosol route.

(continued on next page)

### Further reading, resources and contacts (*continued*)

Perdue, M.L. and Swayne, D.E. (2005). Public health risk from avian influenza viruses. *Avian Diseases* 49, 317–327

Describes the role of the animal–human interface as it pertains to human infection following exposure to avian and mammalian species infected with influenza viruses.

Brankston, G. et al. (2007). Transmission of influenza A in human beings. *Lancet Infectious Diseases* 7, 257–265  
Review of influenza A virus transmission in humans, with emphasis on the contribution of different modes of transmission.

Belser, J.A. et al. (2009). Use of animal models to understand the pandemic potential of highly pathogenic avian influenza viruses. *Advances in Virus Research* 73, 55–97

Describes current animal models for the study of influenza virus pathogenesis and transmission.

### Features associated with this article

#### Figures

Figure 1. Assessing the pandemic potential of influenza viruses.

Figure 2. Contribution of viral molecular determinants in influenza virus transmission.

#### Table

Table 1. Factors influencing behaviour of aerosols in the environment.

### Citation details for this article

Jessica A. Belser, Taronna R. Maines, Terrence M. Tumpey and Jacqueline M. Katz (2010) Influenza A virus transmission: contributing factors and clinical implications. *Expert Rev. Mol. Med.* Vol. 12, e39, December 2010, doi:10.1017/S1462399410001705

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.